

## Original Article

## The Efficacy of Artemether/Lumefantrine for the Treatment of Uncomplicated Falciparum Malaria in Children in an Area of High Transmission in Southern Sudan

Sakina B. Elamin M.Pharm<sup>1</sup>, Gleleldin E. Sidig MD<sup>2</sup>, Khalid A. Elmardi DTM&H<sup>1</sup>, Tarig A. Mohamed MPH<sup>1</sup>, Eldirdiery S. Ahmed MD<sup>3</sup>, Abdelraheem O. Mohamed PhD<sup>4</sup> and Elfatih M. Malik MD<sup>1,5</sup>

<sup>1</sup> National Malaria Control Programme, Khartoum, Sudan, <sup>2</sup> ElSabah Paediatrics Hospital, Juba, Sudan, <sup>3</sup> College of Medicine, University of Juba, Sudan, <sup>4</sup> University of ElImam El Mahdi, Kosti, Sudan, <sup>5</sup> Tel office: +249 183 776809, Tel Home: +249 185 361942, Fax: +249 183 770397, P.O. Box: 1204 Khartoum, e-mail: fatihmmalik@hotmail.com

### Abstract

**Objectives:** Sudanese health authority adopted a new antimalarial drug policy in response to the reported high level of chloroquine resistance. "Artesunate+ sulfadoxine/pyrimethamine" (ASP) and "artemether/lumefantrine" (A/L) are recommended as first and second lines for the treatment of uncomplicated falciparum malaria respectively. This study aims to evaluate the clinical and parasitological response to A/L and to report any side effects related to the drug in children living in high transmission areas.

**Methods:** This evaluation of the clinical and parasitological response to directly observed treatment with 6 doses A/L following WHO protocol for monitoring antimalarial drugs efficacy. Giemsa-stained thick and thin blood smears were examined microscopically on days 0, 2, 3, 7, 14, 21 and 28, or at other times if a patient felt unwell. The primary end point was the 28-day cure rate and the secondary end points were time to parasite clearance.

**Results:** A total of 75 (40.8%) patients met the inclusion criteria, of them 70 (97.2%) patients showed adequate clinical and parasitological response (radical cure), 2 (2.8%) patients were classified as early treatment failure and 3 (4%) patients were lost to be followed-up by day 3. No progression to severe illness or danger signs occurred for any patient during the study.

**Conclusion:** The main outcome of study was that A/L was found to be highly effective against *Plasmodium falciparum* uncomplicated malaria, well tolerated by children with no reported serious side effects. Dramatic decrease in parasites density and fever were observed in most of the cases by day 3 of treatment.